

Atomic few-body systems with muonium

Alexei M. Frolov*

ITAMP, Harvard-Smithsonian Center for Astrophysics,

MS 14, 60 Garden Street, Cambridge MA 02138-1516, USA and

Department of Applied Mathematics

University of Western Ontario, London, Ontario N6H 5B7, Canada

(Dated: January 30, 2017)

Abstract

Properties of some few-body systems which include one positively charged muon μ^+ and two electrons e^- are discussed. In particular, we consider the negatively charged muonium ion Mu^- (or $\mu^+e_2^-$) and four-body MuPs (or $\mu^+e_2^-e^+$) systems each of which has only one stable bound (ground) state. The problem of annihilation of the electron-positron pair(s) in the MuPs system is investigated. The hyperfine structure splitting of the ground state in the MuPs system evaluated with our expectation value of the muon-positron delta-function is $\Delta \approx 23.05758 \text{ MHz}$. Another group of interesting four-body neutral systems investigated in this study includes the $p^+\mu^+e_2^-$, $d^+\mu^+e_2^-$ and $t^+\mu^+e_2^-$ ‘quasi-molecules’. These quasi-molecules are formed in large numbers when positively charged muons slow down in liquid hydrogen, or in liquid deuterium and/or tritium. The properties of these systems are unique, since they occupy an intermediate position between actual two-center molecules and one-center atoms.

PACS number(s): 32.10.Fn, 31.15.A- and 31.15.Ve

*E-mail address: afrolov@uwo.ca

I. INTRODUCTION

In this communication we report the results of our analysis of some three- and four-body systems each of which include one positively charged muon μ^+ and two electron e^- . Briefly, we can say that each of these few-body systems contains muonium Mu (or μ^+e^-), or muonium ion Mu^- (or $\mu^+e_2^-$). Recently, there is an increasing experimental interest to such few-body systems (see discussions and references in [1] - [5]). This can be explained by rapidly growing experimental abilities to detect and isolate similar few-particle systems. Moreover, by using the modern experimental techniques one can investigate some of the bound state properties of these systems. Another reason follows from the fact that the μ^+ muon has positive electric charge and relatively small particle mass. This means that all atomic and molecular few-body systems which contain one positively charged muon and two electrons have very special electron density distribution which differs substantially from electron density distributions in ‘similar’ atomic and molecular systems. In reality, such systems can be considered as a separate class of bound systems which are neither atoms, nor molecules. In some cases, the positively charged muon plays a role of central particle which stabilize a few-electron structure, e.g., in the Mu^- ion and MuPs system (see below). Formally, all few-body systems with positively charged muon are unstable, but their lifetime $\tau (\approx 2 \cdot 10^{-6} \text{ sec})$ significantly exceeds the mean time(s) of atomic transitions and decay processes $\tau_{tr} \leq 1 \cdot 10^{-11} \text{ sec}$. In other words, these few-body quasi-atomic systems can be created and stabilized in their bound state(s) substantially faster than the decay of the μ^+ muon can occur.

As follows from the above, it is important to predict the overall stability of the few-body systems which include muonium Mu (or muonium ion Mu^-) and investigate their basic properties. These problems are considered in our study. In particular, below we consider the negatively charged Mu^- (or $\mu^+e_2^-$) ion and neutral four-body MuPs (or $\mu^+e_2^-e^+$) systems. Each of these systems has only one bound (ground) state. Recently, we could drastically improve the overall accuracy of our variational computations for the both Mu^- ion and MuPs system. For the ground $1S^1$ -state in the Mu^- ion we can now produce the variational wave functions which contain 2 - 3 additional (correct) decimal digits (compare with results from [8]) in the total energy and other bound state properties. The overall accuracy of the bound state computations for the Mu^- ion is comparable with the accuracy of the best

atomic calculations for the two-electron atoms and ions. For the four-body MuPs system we have also derived much better variational approximations to the exact wave function. This allows us to evaluate many bound state properties of this system to very good accuracy (compare with [12]). For the MuPs system we also re-derive the formulas which have been obtained and used earlier [12] for some annihilation rates of the (e^+, e^-) -pair. Finally, a few corrections in such formulas have been made.

A separate group of systems discussed below include three neutral $p^+\mu^+e_2^-$, $d^+\mu^+e_2^-$ and $t^+\mu^+e_2^-$ ‘quasi-molecules’. These four-body systems have never been considered earlier. Note that each of these three systems also contains muonium Mu (or μ^+e^-), or negatively charged muonium ion Mu^- (or $\mu^+e_2^-$). The energy spectra and bound state properties of these four-body systems are substantially different from the spectra of the Mu^- ion and MuPs systems. For instance, the bound state spectrum of each of the $p^+\mu^+e_2^-$, $d^+\mu^+e_2^-$ and $t^+\mu^+e_2^-$ quasi-molecules contains dozens of bound states which converge to the corresponding dissociation threshold(s). These three atomic systems are original, since they cannot be considered as the regular ‘two-center’ molecules even in the first approximation. On the other hand, these systems are not one-center atoms (or quasi-atoms). Formally, these three neutral $p^+\mu^+e_2^-$, $d^+\mu^+e_2^-$ and $t^+\mu^+e_2^-$ systems form a new sub-class among all atomic and molecular systems.

This paper has the following structure. The negatively charged muonium ion Mu^- is considered in the next Section. Since this ion is a three-body system, then we can determine the total energy of the ground 1^1S -state of the Mu^- ion to very high numerical accuracy. Many expectation values (i.e. properties) of this ion in its ground state have also been determined in our highly accurate computations. The neutral four-body MuPs system is considered in the third Section. Here we also evaluate a number of few-photon annihilation rates for this system and the hyperfine structure splitting. The four-body $p^+\mu^+e_2^-$, $d^+\mu^+e_2^-$ and $t^+\mu^+e_2^-$ quasi-molecules are discussed in the fourth Section. Note that these systems have never been considered in earlier studies. For these three systems we consider only their electron ground states. Concluding remarks can be found in the last Section.

II. NEGATIVELY CHARGED MUONIUM ION

First, let us consider the negatively charged three-body muonium ion Mu^- which consists of one positively charged muon μ^+ and two electrons e^- . As mentioned above this system has only one stable bound state which is the ground 1^1S –electron state. Stability of the ground state in the Mu^- ion has been shown in 1986 [6]. A few years later this ion has been created in actual experiment [7]. The total energies and other bound state properties of this ion have been considered in [8]. Since our paper [8] was published the new experiments in high energy physics with positively charged muons have been conducted. The improved numerical value of the muonic mass now is:

$$m_\mu = \frac{105.65836668 \text{ MeV}/c^2}{0.510998910 \text{ MeV}/c^2} = 206.76828191 \dots m_e \quad (1)$$

The non-relativistic Hamiltonian of the Mu^- ion takes the form

$$H = -\frac{\hbar^2}{2m_e} \left[\nabla_1^2 + \nabla_2^2 + \frac{m_e}{m_\mu} \nabla_3^2 \right] - \frac{e^2}{r_{32}} - \frac{e^2}{r_{31}} + \frac{e^2}{r_{21}} \quad , \quad (2)$$

where $\hbar = \frac{h}{2\pi}$ is the reduced Planck constant, m_e is the electron mass and e is the absolute value of the electric charge of electron. In this equation the subscripts 1 and 2 designate the two electrons e^- , while the subscript 3 denotes the heavy central particle μ^+ (quasi-nucleus) with the mass m_μ ($m_\mu \gg m_e$), and positive electric (nuclear) charge $+e$. For the Mu^- ion we need to solve the non-relativistic Schrödinger equation $H\Psi = E\Psi$ for the two-electron ions, where H is the Hamiltonian, Eq.(2), and E ($E < 0$) is the eigenvalue, which coincides with the total energy of the Mu^- ion in its ground 1^1S –state. This state will be stable, if its total energy is lower than the total energy of the two-body muonium Mu (μ^+e^-) which equals $-\frac{m_\mu}{2(m_\mu+1)} = 0.5(1 + m_\mu^{-1})$ in atomic units.

Numerical solution of the Schrödinger equation allows one to obtain highly accurate wave function(s) Ψ . In reality, the unknown (exact) wave function is approximated by using some rapidly convergent variational expansions. One of the best variational expansions known for three-body systems is the exponential variational expansion in the relative coordinates r_{32}, r_{31}, r_{21} , or in the perimetric coordinates u_1, u_2, u_3 . For the ground (bound) 1^1S –state of the two-electron ions/atoms the explicit form of these expansions are

$$\Psi = \frac{1}{2}(1 + \hat{P}_{12}) \sum_{i=1}^N C_i \exp(-\alpha_i r_{32} - \beta_i r_{31} - \gamma_i r_{21})$$

$$= \frac{1}{2}(1 + \hat{P}_{12}) \sum_{i=1}^N C_i \exp[-(\alpha_i + \beta_i)u_3 - (\alpha_i + \gamma_i)u_2 - (\beta_i + \gamma_i)u_3] \quad (3)$$

where $r_{ij} = |\mathbf{r}_i - \mathbf{r}_j| = r_{ji}$ are the interparticle scalar distances and $\mathbf{r}_i$ ($i = 1, 2, 3$) are the three Cartesian coordinates of the particles. In Eq.(3) the notation $\hat{P}_{12}$ stands for the permutation operator of identical particles (electrons), C_i ($i = 1, 2, \dots, N$) are the linear parameters of the exponential expansion, Eq.(3), while α_i, β_i and γ_i are the non-linear parameters of this expansion. These non-linear parameters must be varied in calculations to increase the overall efficiency and accuracy of the method. The best-to-date optimization strategy for these non-linear parameters was described in [9], while its modified (advanced) version was presented in another paper published in 2006 (see the second paper in Ref.[9]).

Note that the $3N$ following conditions $\alpha_i + \beta_i > 0, \alpha_i + \gamma_i > 0, \beta_i + \gamma_i > 0$ (for $i = 1, 2, \dots, N$) must always be obeyed for the non-linear parameters in Eq.(3) to guarantee convergence of all three-particle integrals which are needed in computations. Three scalar perimetric coordinates u_i are simply related with the relative coordinates r_{kj} by the following linear relations

$$\begin{aligned} u_1 &= \frac{1}{2}(r_{21} + r_{31} - r_{32}) \quad , \quad r_{32} = u_2 + u_3 \\ u_2 &= \frac{1}{2}(r_{21} + r_{32} - r_{31}) \quad , \quad r_{31} = u_1 + u_3 \\ u_3 &= \frac{1}{2}(r_{31} + r_{32} - r_{21}) \quad , \quad r_{21} = u_1 + u_2 \end{aligned} \quad (4)$$

where $r_{ij} = r_{ji}$. In contrast with the relative coordinates r_{32}, r_{31}, r_{21} the three perimetric coordinates u_1, u_2, u_3 are independent of each other and each of them varies between 0 and $+\infty$. This drastically simplifies analytical and numerical computations of all three-body integrals which are needed for solution of the corresponding eigenvalue problem and for evaluation a large number of bound state properties of the Mu^- ion. Additional interesting details about perimetric coordinates can be found in the Appendix.

The total energies and bound state properties of this ion (in atomic units) can be found in Tables I (energies) and II (properties). All these values have been determined to very high numerical accuracy with the use of the extended precision arithmetic [10]. Analogous bound state properties of the Mu^- ion are slightly different from values presented in [8] which can be explained by the differences in the muonium mass m_μ . It should be mentioned that the current accuracy of numerical computations for three-particle (non-relativistic) Coulomb

systems is extremely high and this allows one to determine the energies and wave functions of all bound states to very high accuracy which can be evaluated as $\approx 1 \cdot 10^{-22} - 1 \cdot 10^{-25}$ *a.u.* The overall accuracy of our numerical calculations of the ground 1^1S -state in the Mu^- ion $\approx 3 \cdot 10^{-24}$ *a.u.* is quite comparable to the accuracy of the best atomic bound state computations for two-electron atoms and ions. Highly accurate wave functions can be used to obtain the correct numerical values of many bound state properties, including a few slowly convergent properties [11], e.g., the $\langle r_{ij}^6 \rangle$, $\langle r_{ij}^8 \rangle$ and $\langle r_{ij}^{10} \rangle$ expectation values. As expected the numerical values of all computed bound state properties of the Mu^- ion are close to the analogous properties of the negatively charged hydrogen ion H^- which are also shown in Table II (see [12]). The results from Table I and II is the most complete and accurate set of the expectation values ever determined for the muonium ion Mu^- .

The negatively charged muon Mu^- ion is an example of three-body quasi-atomic system where the μ^+ muon plays a role of a central particle, or ‘quasi-nucleus’. Another such a system with the same quasi-nucleus is the four-body neutral system MuPs (or $\mu^+e_2^-e^+$) which is discussed below. Note that the MuPs system has a more complicated internal structure and a number of interesting bound state properties.

III. FOUR-BODY MUONIUM-POSITRONIUM SYSTEM

Another systems which contains muonium Mu (μ^+e^-), or muonium ion Mu^- ($\mu^+e_2^-$) is the four-body MuPs (or $\mu^+e_2^-e^+$) quasi-atom which contains one bound positron [12]. In other words, the MuPs ‘quasi-atom’ is a pure leptonic four-body quasi-atom which also contains one positively charged muon (heavy ‘central’ particle), two electrons and one positrons (light particles). Such a unique system can now be created and observed in actual experiments in which a large number of positively charged muons are created. The stability of the MuPs system follows from the fact that the analogous four-body HPs and Ps_2 systems are stable. Moreover, since each of the HPs and Ps_2 systems has only one bound state (ground state), then we conclude that the analogous MuPs systems also has only one bound (ground) state which is the 1^1S -electron state. In atomic units the non-relativistic Hamiltonian of the four-body MuPs system takes the form (see, e.g., [13])

$$H = -\frac{1}{2} \left[\nabla_1^2 + \nabla_2^2 + \nabla_3^2 + \frac{1}{m_\mu} \nabla_4^2 \right] - \frac{1}{r_{32}} - \frac{1}{r_{31}} - \frac{1}{r_{42}} - \frac{1}{r_{41}} + \frac{1}{r_{21}} + \frac{1}{r_{43}} \quad , \quad (5)$$

where the particles 1 and 2 are the two electrons e^- (or ‘-’), while the particles 3 and 4 are the positron e^+ (or ‘+’) and muon μ^+ , respectively. The MuPs system contains two light negative particles and one light positive particle positron e^+ which are moving around one heavy center μ^+ . In this sense the MuPs system is similar to the four-body HPs system(s). The mass of the positively charged muon used in this Section is $m_\mu = 206.768264m_e$.

To determine the accurate solution(s) of the non-relativistic Schrödinger equation $H\Psi = E\Psi$ in this Section we apply the variational expansion of the wave function written in multi-dimensional gaussoids, which are, in fact, the six-dimensional gaussoids for the four-body MuPs system. Each of the basis functions in this expansion explicitly depends upon all six relative coordinates r_{ij} , where $r_{ij} = r_{12}, r_{13}, r_{14}, r_{23}, r_{24}$ and r_{34} . For the ground state of the MuPs system the variational expansion in six-dimensional gaussoids takes the form (see, e.g., [14], [15]):

$$\psi(L=0; S=0) = \frac{1}{2}(1 + \hat{P}_{12}) \sum_{i=1}^{N_A} C_i \exp(-a_{ij}r_{ij}^2) \chi_{S=0} \quad (6)$$

where C_i are the linear variational coefficients of the variational function, while a_{ij} , where $(ij) = (12), (13), \dots, (34)$, are the six non-linear parameters in the radial function associated with the $\chi_{S=0}$ spin function. This spin function can be chosen in the form

$$\chi_{S=0} = (\alpha_1\beta_2 - \beta_1\alpha_2)\alpha_\mu\alpha_+ = \alpha_4\alpha_3(\alpha_1\beta_2 - \beta_1\alpha_2) \quad (7)$$

where the notations α and β are used spin-up and spin-down functions (see, e.g., [13]). The total energies and bound state properties of the MuPs system (in atomic units) determined with the use of the trial functions, Eq.(6), can be found in Table III. In general, the current numerical computations of the bound states in the four-body systems with comparable particle masses is significantly less than such an accuracy obtained in calculations of the Coulomb three-body systems (see, e.g., the Mu^- ion discussed above). This can be explained by a different differential geometry of the general four-body problem. Briefly, we can say that follows from the fact that it is impossible to define six perimetric coordinates (for four-body systems) which have the same properties as the three perimetric coordinates introduced above for three-body systems (see Appendix).

The expectation values from Table III can be considered as ‘elementary’ properties. However, by using these ‘elementary’ properties we can determine (or evaluate) some actual properties of the MuPs system. In particular, by using the expectation value of the

electron-positron delta-function from Table III we can evaluate the two- and three-photon annihilation rates $\Gamma_{2\gamma}$ and $\Gamma_{3\gamma}$ for the ground state in the MuPs system. The sum of these two annihilation rates essentially determines the life-time of the MuPs system. The formula for the $\Gamma_{2\gamma}$ rate takes the form [12]

$$\Gamma_{2\gamma}(\text{MuPs}) = 2\pi\alpha^4 c a_0^{-1} \langle \delta(\mathbf{r}_{+-}) \rangle = 100.939408683 \cdot 10^9 \langle \delta(\mathbf{r}_{+-}) \rangle \text{ sec}^{-1} \quad (8)$$

where $\langle \delta_{+-} \rangle$ is the expectation value of the electron-positron delta-function determined for the ground (bound) 1^1S_e -state in the MuPs system. The indexes '+' and '-' used in delta-functions in Eq.(8) designate the positron and electron, respectively, while the index μ stands for the μ^+ muon. In Eq.(8) and everywhere below the notation $\alpha = \frac{e^2}{\hbar c} = 7.2973525698 \cdot 10^{-3} \left(\approx \frac{1}{137} \right)$ is the dimensionless fine-structure constant, $c = 2.99792458 \cdot 10^8 \text{ m} \cdot \text{sec}^{-1}$ is the speed of light in vacuum, and a_0 is the Bohr radius which equals $0.52917721092 \cdot 10^{-10} \text{ m}$ [16]. Analogous formula for the three-photon annihilation rate $\Gamma_{3\gamma}(\text{MuPs})$ takes the form

$$\Gamma_{3\gamma}(\text{MuPs}) = 2 \frac{4(\pi^2 - 9)}{3} \alpha^5 c a_0^{-1} \langle \delta(\mathbf{r}_{+-}) \rangle = 2.718545954 \cdot 10^8 \langle \delta(\mathbf{r}_{+-}) \rangle \text{ sec}^{-1} \quad (9)$$

This formula can easily be derived from the well known formula obtained in [17] for one triplet electron-positron pair. The lowest-order QED correction to the $\Gamma_{2\gamma}(\text{MuPs})$ annihilation rate is written in the form (derived from the formula [18] for one singlet electron-positron pair)

$$\Delta\Gamma_{2\gamma}(\text{MuPs}) = -2\pi\alpha^5 c a_0^{-1} \left(\frac{13\pi}{12} - \frac{7}{\pi} \right) \langle \delta(\mathbf{r}_{+-}) \rangle = -8.6565792289 \cdot 10^8 \langle \delta(\mathbf{r}_{+-}) \rangle \text{ sec}^{-1} \quad (10)$$

This correction is negative. By using these formulas and our expectation value for the electron-positron delta-function $\langle \delta(\mathbf{r}_{+-}) \rangle$ from Table III we have found that $\Gamma_{2\gamma}(\text{MuPs}) = 2.464793493 \cdot 10^9 \text{ sec}^{-1}$, $\Gamma_{3\gamma}(\text{MuPs}) = 6.63829367 \cdot 10^6 \text{ sec}^{-1}$ and $\Delta\Gamma_{2\gamma}(\text{MuPs}) = -2.1138106945 \cdot 10^7 \text{ sec}^{-1}$, respectively. Numerical values of the four- and five-photon annihilation rates ($\Gamma_{4\gamma}$ and $\Gamma_{5\gamma}$) are obtained from the $\Gamma_{2\gamma}(\text{MuPs})$ and $\Gamma_{3\gamma}(\text{MuPs})$ rates mentioned above with the use of the following formulas (derived from the original formulas obtained in [19])

$$\Gamma_{4\gamma}(\text{MuPs}) = 0.274 \left(\frac{\alpha}{\pi} \right)^2 \Gamma_{2\gamma}(\text{MuPs}) \quad , \quad \Gamma_{5\gamma}(\text{MuPs}) = 0.177 \left(\frac{\alpha}{\pi} \right)^2 \Gamma_{3\gamma}(\text{MuPs}) \quad . \quad (11)$$

With our expectation values of the electron-positron delta-function $\delta(\mathbf{r}_{+-})$ (see Table III) are $3.64386 \cdot 10^3 \text{ sec}^{-1}$ and 6.33958 sec^{-1} .

The one-photon positron annihilation in the MuPs system is of interest in some problems as well as for the future development of QED. This process can be represented as a regular

two-photon annihilation of the (e^-, e^+) -pair and the following absorbtion of one of the emitted photons either by the second electron, or by the μ^+ muon (heavy particle). Therefore, one can observe, in principle, the two one-photon annihilations in MuPs. The rates of these processes are designated below as $\Gamma_{1\gamma}^{(e)}(\text{MuPs})$ and $\Gamma_{1\gamma}^{(\mu)}(\text{MuPs})$. Analytical expressions for these rates can be derived from the results of earlier studies [20] and [21]. In atomic units the explicit formulas for the $\Gamma_{1\gamma}^{(e)}(\text{MuPs})$ and $\Gamma_{1\gamma}^{(\mu)}(\text{MuPs})$ annihilation rates take the form

$$\Gamma_{1\gamma}^{(e)}(\text{MuPs}) = \frac{64\pi^2}{27}\alpha^8 c a_0^{-1} \langle \delta(\mathbf{r}_{+--}) \rangle = 1065.75691955 \langle \delta(\mathbf{r}_{+--}) \rangle \quad , \quad (12)$$

and

$$\Gamma_{1\gamma}^{(\mu)}(\text{MuPs}) = \frac{4\pi^2}{3}\alpha^8 Q^5 c a_0^{-1} \langle \delta(\mathbf{r}_{+-\mu}) \rangle = 599.48826725 \langle \delta(\mathbf{r}_{+-\mu}) \rangle \quad , \quad (13)$$

where $Q = 1$ for MuPs, while $\langle \delta(\mathbf{r}_{+--}) \rangle$ and $\langle \delta(\mathbf{r}_{+-\mu}) \rangle$ are the expectation values of the corresponding triple delta-functions determined in atomic units. In derivation of the formula, Eq.(13), we have assumed that the muonium mass is very large (infinite). The sum of these values allows one to determine the one-photon annihilation rate for the MuPs system

$$\Gamma_{1\gamma}(\text{MuPs}) = \frac{4\pi^2}{3}\alpha^8 c a_0^{-1} \left[\langle \delta(\mathbf{r}_{+-\mu}) \rangle + \frac{16}{9} \langle \delta(\mathbf{r}_{+--}) \rangle \right] \quad , \quad (14)$$

By using our expectation values of the triple delta-functions mentioned in these equations (see Table III) we have found the following numerical values $\Gamma_{1\gamma}^{(e)}(\text{MuPs}) \approx 0.39087487 \text{ sec}^{-1}$, $\Gamma_{1\gamma}^{(\mu)}(\text{MuPs}) \approx 0.51148759 \text{ sec}^{-1}$ and $\Gamma_{1\gamma}(\text{MuPs}) \approx 0.90236246 \text{ sec}^{-1}$.

We can also evaluate the hyperfine structure splitting in the MuPs system, where the hyperfine structure arises from direct interaction between the spin-vectors of the positron and muon. The hyperfine structure splitting in the MuPs system $(\Delta H)_{hsp}$ is written in the form [12]

$$(\Delta H)_{hsp} = \frac{8\pi\alpha^2}{3} \mu_B^2 \frac{g_\mu}{m_\mu} \frac{g_e}{m_e} \langle \delta(\mathbf{r}_{+\mu}) \rangle = 14229.1255 \langle \delta(\mathbf{r}_{+\mu}) \rangle \quad (15)$$

where α is the fine-structure constant, μ_B is the Bohr magneton which equals 0.5 in atomic units, while $\langle \delta_{+\mu} \rangle$ is the expectation value of the muon-positron delta-function. Also in Eq.(15) the notations m_μ and m_e stand for the muon/positron masses-at-rest, while the factors $g_+ = -2.0023193043718$ and $g_\mu = -2.0023318396$ are the gyromagnetic ratios. In Eq.(15) we have used the factor $6.579 \ 683 \ 920 \ 61 \cdot 10^9 \text{ (MHz/a.u.)}$ has been used to recalculate the ΔE_{hf} energy from atomic units to *MegaHertz*. By using the expectation value of the muon-positron delta-function $\langle \delta_{+\mu} \rangle \approx 1.62045 \cdot 10^{-3} \text{ a.u.}$ from Table III, one

finds that the value a in Eq.(15) equals $(\Delta H)_{hsp} \approx 23.05758 \text{ MHz}$. This coincides with the energy difference between the hyperfine structure states with $J = 0$ and $J = 1$, where the notation J stands for the total spin of the muon-positron pair in the MuPs system.

To conclude this Section let us note that there is another interesting process which can be observed, in principle, in the MuPs system. This process is the conversion of the $\mu^+ - e^-$ pair into its charge-conjugate $\mu^- - e^+$ pair. During the original MuPs system is transformed into another (new) Mu*Ps system which is also bound and contains two positrons. The overall probability of this process is very small, but it is proportional to the $\langle \delta(\mathbf{r}_{-\mu}) \rangle$ expectation value (see Table III).

IV. FOUR-BODY HYDROGEN-MUONIUM QUASI-MOLECULES

The third group of few-body systems with positively charged muons considered in this study includes three four-body ‘quasi-molecules’: $p^+\mu^+e_2^-$, $d^+\mu^+e_2^-$ and $t^+\mu^+e_2^-$. These systems are always formed when high-energy positively charged muons slow down in the liquid hydrogen, or in liquid deuterium and/or tritium. Each of these quasi-molecules includes one positively charged muon and two bound electrons. In each of these three quasi-molecules the μ^+ muon plays the role of the second positively charged ‘heavy’ center. Indeed, as follows from the mass ratio of the muon and electron masses the muon’s velocity in the $a\mu e_2$ ($= a^+\mu^+e_2^-$) system is substantially smaller (~ 206.768 times smaller) than the electron’s velocity. This means that in the first approximation the μ^+ muon can be considered as the second heavy particle in the $a\mu e_2$ quasi-molecule, where $a = p, d, t$. However, the actual small dimensionless parameters in the Born-Oppenheimer theory [22] (see also [23]) of the adiabatic two-center systems are defined as follows (in our notations):

$$\tau_1 = \sqrt[4]{\frac{M_a}{M_a + m_\mu + 2m_e}} \quad \text{and} \quad \tau_2 = \sqrt[4]{\frac{m_\mu}{M_a + m_\mu + 2m_e}} \quad (16)$$

where $a = (p, d, t)$ and notations m_μ and M_a stand for the masses of the two heavy particles. In our calculations we used the following numerical values for the particle masses: $M_p = 1836.152701m_e$, $M_d = 3670.483014m_e$, $M_t = 5496.92158m_e$ and $m_\mu = 206.768262m_e$. The two dimensionless parameters τ_1 and τ_2 determine those systems to which adiabatic approximation can be applied. Formally, the adiabatic approximation works well in those cases when $\tau_{min} = \min(\tau_1, \tau_2) \geq 0.5$. For the $p^+\mu^+e_2^-$, $d^+\mu^+e_2^-$ and $t^+\mu^+e_2^-$ quasi-molecules

the parameter τ_{min} does not exceed 0.15, i.e. it is significantly smaller than 0.5. This means that any of the three $a\mu e_2$ systems investigated in this study cannot be considered as an actual molecule with the two heavy centers (or two-center molecule). The main reason for this is obvious, since the directions of the central $a^+ \leftrightarrow \mu^+$ axis is not fixed in space even in the $t\mu e_2$ system. In other words, the mass of the positively charged μ^+ ‘center’ is too small to consider this particle as an actual ‘second’ heavy center in the $a\mu e_2$ system. In reality, the $p^+\mu^+e_2^-$, $d^+\mu^+e_2^-$ and $t^+\mu^+e_2^-$ quasi-molecules form a separate class of quasi-atomic systems which differs substantially from regular atoms and molecules. Below, we consider these three four-body systems as ‘quasi-molecules’, but this name is confusing, since it does not represent the actual situation with these systems.

In atomic units the non-relativistic Hamiltonian of the four-body $a\mu e_2$ systems is written in the form

$$H = -\frac{1}{2}\left[\nabla_1^2 + \nabla_2^2 + \frac{1}{m_\mu}\nabla_3^2 + \frac{1}{M_a}\nabla_4^2\right] - \frac{e^2}{r_{32}} - \frac{e^2}{r_{31}} - \frac{e^2}{r_{42}} - \frac{e^2}{r_{41}} + \frac{e^2}{r_{21}} + \frac{e^2}{r_{43}}, \quad (17)$$

where $e = 1$ and the particles 1 and 2 are the two electrons, while the particles 3 and 4 are the positively charged muon and heavy particle a , where $a = p, d, t$, respectively. In Eq.(17) we assume that the masses of two heavy particles a^+ and μ^+ are expressed in terms of the electron mass m_e . The $a\mu e_2$ system contains two light negative particles (electrons) and two heavy positive particles (or two centers) which form the ‘molecular’ axis $a^+ \leftrightarrow \mu^+$. For pure adiabatic two-center molecule the location of this axis in space must be fixed, but this is not the case for the $p^+\mu^+e_2^-$, $d^+\mu^+e_2^-$ and $t^+\mu^+e_2^-$ quasi-molecules.

In general, the properties of the $a\mu e_2$ quasi-molecular system must be similar to the properties of the two-center hydrogen molecules such as D_2 , DT and/or HD . Direct comparison of the numerical values of bound state properties determined for the $p^+\mu^+e_2^-$, $d^+\mu^+e_2^-$ and $t^+\mu^+e_2^-$ quasi-molecules with the analogous properties of the H_2 molecule indicates clearly that such a similarity is observed. However, the overall accuracy of such a similarity is not high and cannot be improved, e.g., by using more accurate wave functions. In reality, the $p\mu e_2$, $d\mu e_2$ and $t\mu e_2$ quasi-molecular systems form a separate class among four-body systems with two electrons (see above). The same conclusion has been mentioned in [8], where we discussed the analogous three-body ions: $(p\mu e)^+$, $(d\mu e)^+$ and $(t\mu e)^+$ (see also [24] and [25]).

In order to solve the non-relativistic Schrödinger equation $H\Psi = E\Psi$ for the $a^+\mu^+e_2^-$ systems we applied the same variational expansion in multi-dimensional gaussoids, Eq.(6). The

non-linear parameters a_{ij} in this expansion, Eq(6), must carefully be optimized to produce accurate numerical results, including the total energies of these systems. By performing a number of consecutive optimizations of these parameters we have produced a number of trial wave functions which can be considered as accurate approximations to the actual wave functions of these four-body systems. A large number of bound state properties of the $p^+\mu^+e_2^-, d^+\mu^+e_2^-$ and $t^+\mu^+e_2^-$ quasi-molecules in their ground state(s) can be found in Table III. By using the expectation values from Table III one can predict numerical values of some actual properties, i.e properties which can be observed in experiments. For instance, by using the expectation values of the muon-proton (or muon-deuteron, etc) delta-function ($\langle\delta(\mathbf{r}_{a\mu})\rangle$ in our notations) we can evaluate the hyperfine structure splitting in each of the $a\mu e_2$ considered in this study.

The hyperfine structure splitting in the ground S -states of the four-body $p^+\mu^+e_2^-, d^+\mu^+e_2^-$ and $t^+\mu^+e_2^-$ quasi-molecules is the result of the direct spin-spin interaction between muonic spin and spin of the heavy hydrogen nucleus, i.e. protium, deuterium and tritium. The two electrons form a singlet pair with zero contribution into the hyperfine structure. The formula which is used to evaluate the hyperfine structure splitting $(\Delta H)_{hsp}$ in the ground S -states of each of the four-body $a\mu e_2$ quasi-molecule takes the form (in atomic units)

$$(\Delta H)_{hsp} = \frac{2\pi}{3}\alpha^2 \frac{g_N g_\mu}{m_p m_\mu} \langle\delta(\mathbf{r}_{\mu N})\rangle (\mathbf{I}_N \cdot \mathbf{s}_\mu) = \frac{\pi}{3}\alpha^2 \frac{g_N g_\mu}{m_p m_\mu} \langle\delta(\mathbf{r}_{\mu N})\rangle [J(J+1) - I_N(I_N+1) - s_\mu(s_\mu+1)] \quad , \quad (18)$$

where $\langle\delta(\mathbf{r}_{\mu N})\rangle$ is the expectation value of the muon-nucleus delta-function, while other notations have exactly the same meaning and numerical values as in Eq.(15). Note that in atomic units the Bohr magneton μ_B exactly equals $\frac{1}{2}$. The factor g_N in Eq.(15) equals to the ratio $\frac{\mathcal{M}}{I_N}$, where $\mathcal{M}_p = 2.792847386$, $\mathcal{M}_d = 0.857438230$ and $\mathcal{M}_t = 2.9789624775$, while $I_p = \frac{1}{2}$, $I_d = 1$ and $I_t = \frac{1}{2}$. The factor $m_p = 1836.152701 m_e$ is the proton's mass expressed in the electron mass m_e . Also, in Eq.(18) the factor $6.579\,683\,920\,61 \cdot 10^9$ (MHz/a.u.) must be used to re-calculate the ΔE_{hf} energy from atomic units to *MegaHertz*. By using these numerical values we can evaluate the coefficients which arise in Eq.(18) for the $p\mu e_2, d\mu e_2$ and $t\mu e_2$ quasi-molecules. These coefficients are: -10.808930550, -1.651657610 and -11.529236682, respectively. Now, by using the expectation values of the muon-nucleus delta-functions from Table III we obtain the following hyperfine structure for these four-body

systems: (a) $\varepsilon(J = 1) = -30.098 \text{ Hz}$ and $\varepsilon(J = 0) = 90.295 \text{ Hz}$ for $p\mu e_2$, (b) $\varepsilon(J = \frac{3}{2}) = -12.567 \text{ Hz}$ and $\varepsilon(J = \frac{1}{2}) = 25.135 \text{ Hz}$ for $d\mu e_2$, and (c) $\varepsilon(J = 1) = -46.023 \text{ Hz}$ and $\varepsilon(J = 0) = 138.070 \text{ Hz}$ for $t\mu e_2$. The differences between these two energies gives the corresponding hyperfine structure splitting for each quasi-molecule (120.393 Hz, 37.702 Hz and 184.093 Hz, respectively). Very small values of these hyperfine structure splittings indicate that each of these system has internal structure which is close to the adiabatic two-center molecule. On the other hand, these numerical values are in dozens times larger than values which can be found in any actual two center molecule.

V. CONCLUSION

We have considered a number of systems which contain the muonium Mu, or muonium ion Mu^- . In other words, these systems include one positively charged muon μ^+ and two boound electrons e^- . The total energies and a large number of bound state properties of the negatively charged muonium ion Mu^- have been obtained from highly accurate numerical computations. The ground state in the four-body MuPs (or $\mu^+e_2^-e^+$) system is also studied in detail. It is shown that the bound state properties are similar to analogous properties of the HPs four-body system. For MuPs we evaluate the two- and three-photon annihilation rates and hyperfine structure splitting. The computed expectation values agree very well with the results of earlier studies (see, e.g., [12]), but our current results have significantly better numerical accuracy.

In this study we also investigate the bound (ground) states in the four-body quasi-molecules $p^+\mu^+e_2^-$, $d^+\mu^+e_2^-$ and $t^+\mu^+e_2^-$ (or $p\mu e_2$, $d\mu e_2$ and $t\mu e_2$). These systems have never been considered in earlier studies. The three quasi-molecules $p\mu e_2$, $d\mu e_2$ and $t\mu e_2$ cannot be considered as the ‘pure adiabatic’ two-center molecules, since the positively charged muon μ^+ is a relatively light particle. This fact allows us to apply some regular methods, e.g., the variational expansion, Eq.(6), to study the internal structure of these ‘intermediate’ systems. If the $p^+\mu^+e_2^-$, $d^+\mu^+e_2^-$ and $t^+\mu^+e_2^-$ would be truly adiabatic (or two-center) systems, then the expansion, Eq.(6), could not be used directly, i.e. without modifications, to produce highly accurate results for such systems. By using our ‘regular’ variational expansion in six-dimensional gaussoids we have determined a number of bound state properties of the ground state(s) in these four-body quasi-molecules. In addition to this we investigated the hyperfine

structure and evaluated the hyperfine structure splitting in each of the $p^+\mu^+e_2^-$, $d^+\mu^+e_2^-$ and $t^+\mu^+e_2^-$ systems.

VI. APPENDIX. PERIMETRIC COORDINATES

Note that the three-body perimetric coordinates u_1, u_2, u_3 were known to antic greeks. The Heron's formula (derived first by Hero of Alexandria, but very likely it was also known to Archimedes 200 years earlier) gives the area of a triangle S by whose sides have lengths r_{32}, r_{31} and r_{21} :

$$S = \sqrt{pu_1u_2u_3} \quad (19)$$

where u_1, u_2, u_3 are the perimetric coordinates defined above (see, Eq.(4)), while $p = \frac{1}{2}(u_1 + u_2 + u_3) = \frac{1}{2}(r_{32} + r_{31} + r_{21})$ is the semi-perimeter of this triangle which coincides with the half of the sum of three perimetric coordinates (and three relative coordinates). Here we assume that the indexes 1, 2 and 3 stand for the three vertexes of the triangle. To transform this formula into the actual three-body problem we need to place three point particles in three vertexes of of this triangle. Simplicity of the formula, Eq.(19), is the first indication of high efficiency of the perimetric coordinates in application to the analysis of various three-body problems. Three perimetric coordinates have been introduced in actual numerical computations of three-body systems by C.L. Pekeris almost 60 years ago [26], [27]. The use of these coordinates was crucial to obtain highly accurate solutions for a large number of three-body problems. It is interesting to note that C.L. Pekeris always considered himself as a pure 'geophysicist' which has a restricted interest in atomic physics (a 'personal hobby' - according to Yu.N. Demkov who met with Pekeris in the early 1970's). Now a large mehods based on the use of the perimetric coordinates are extensively used in various three-body problems and, in particular, for analytical and numerical computations of different three-body integrals, including very complex and singular integrals.

An obvious success of the perimetric coordinates for various three-body problems stimulated discussion about the four-body perimetric coordinates which can be used for highly accurate solutions of different four-body problems [28]. Indeed, for an arbitrary four-body system we can introduce twelve perimetric coordinates which must obey six additional conditions (or constraints) [28]. Formally, we can exclude six (of twelve) perimetric coordinates

by solving these six constraints as algebraic equations. The six remaining perimetric coordinates will be sufficient for a complete description of an arbitrary four-body problem. However, during such a procedure some useful properties of the perimetric coordinates can be lost, e.g., some of them can be negative, or vary between the lower and upper limits A and B , where $A \neq 0$ and $B \neq +\infty$. In addition to this, we have a problem of re-ordering of the arising perimetric coordinates (this problem does not exist for three-body systems). This drastically complicates analytical and numerical computations of actual four-body systems with the use of the four-body perimetric coordinates. Note also that the problem of the correct definition of the four-body perimetric coordinates has been re-investigated recently [29].

VII. ACKNOWLEDGMENTS

This work was supported in part by the NSF through a grant for the Institute for Theoretical Atomic, Molecular, and Optical Physics (ITAMP) at Harvard University and the Smithsonian Astrophysical Observatory. Also, I wish to thank James Babb (ITAMP) and David M. Wardlaw for stimulating discussion.

-
- [1] *Muonic Atoms and Molecules*, (Eds. L.A. Schaller and C. Petitjean, Monte Verita at Birkhauser, (1993)).
 - [2] D.C. Walker, *Muon and Muonium Chemistry*, (Cambridge University Press, New York (2008)).
 - [3] P. Percival, *Muonium*, Lecture 9, TRIUMF Summer Institute (2011), unpublished.
 - [4] S.F.J. Cox, J.S. Lord, I. McKenzie, J.-J. Adjizian, M.I. Heggie, U.A. Jayasooriya, R. Grinter and I.D. Reid, *J. Phys. Condens. Matter* **23**, 315801 (2011).
 - [5] K.S. Khaw, A. Antognini, T. Prokscha, K. Kirch, L. Liskay, Z. Salman, and P. Crivelli, *Phys. Rev. A* **94**, 022716 (2016).
 - [6] A.M. Frolov, *Zeits. für Phys. D* **2**, 61 (1986).
 - [7] Y. Kuang, K.-P. Arnold, F. Chmely, M. Eckhause, V.W. Hughes, J.R. Kane, S. Kettell, D.-H. Kim, K. Kumar, D.C. Lu, B. Matthias, B. Ni, H. Orth, G. zu Putlitz, H.R. Schaefer, P.A.

- Souder, and K. Woodle, Phys. Rev. A **39**, 6109 (1989).
- [8] A.M. Frolov, Phys. Rev. A **69**, 022505 (2004).
 - [9] A.M. Frolov, Phys. Rev. E **64**, 036704 (2001); *ibid*, **74**, 027702 (2006).
 - [10] D.H. Bailey, AMC Trans. Math. Soft. **21**, 379 (1995).
 - [11] A.M. Frolov and D.M. Wardlaw, Europ. Phys. Journ. B **89**, 190 (2016).
 - [12] A.M. Frolov, Europ. Phys. Journ. D **69**, 50 (2015).
 - [13] L.D. Landau and E.M. Lifshitz, *Quantum Mechanics: Non-Relativistic Theory*, (3rd. ed. Pergamon Press, New York (1976)).
 - [14] N.N. Kolesnikov and V.I. Tarasov, Yad. Phys. **35**, 609 (1982) [Sov. Phys. Nucl. Phys. **35**, 354 (1982)].
 - [15] A.M. Frolov and D.M. Wardlaw, JETP **138**, 5 (2010).
 - [16] *CRC Handbook of Chemistry and Physics*, (95th Edition, Ed. W.M. Haynes, Taylor and Francis Group, Boca Raton, FL, (2014-2015)).
 - [17] A. Ore and I. Powell, Phys. Rev. **75**, 1696 (1949).
 - [18] I. Harris and L.M. Brown, Phys. Rev. **105**, 1656 (1957).
 - [19] G. P. Lepage, P.B. Mackenzie, K. H. Streng, and P. M. Zerwas, Phys. Rev. A **28**, 3090 (1983).
 - [20] S.I. Kryuchkov, J. Phys. B **27**, L61 (1994).
 - [21] E. Fermi and G.E. Uhlenbeck, Phys. Rev. **44**, 510 (1933).
 - [22] M. Born and E. Oppenheimer, Ann. der Physik **84**, 457 (1927).
 - [23] H.A. Bethe, *Intermediate Quantum Mechanics*, (W.A. Benjamin Inc., New York (1964)), Chapt. 11.
 - [24] D. McKenna and B. Webster, J. Chem. Soc., Faraday Trans. **80**, 589 (1984).
 - [25] D. McKenna and B. Webster, J. Chem. Soc., Faraday Trans. **81**, 225 (1985).
 - [26] C.L. Pekeris, Phys. Rev. **112**, 1649 (1958).
 - [27] C.L. Pekeris, Phys. Rev. **115**, 1216 (1959).
 - [28] A.M. Frolov, J. Phys. A **39**, 15421 (2006).
 - [29] M. Kreuter, *The Pekeris Method for Lithium: Possibilities and Obstructions*, Thesis, The University of Wisconsin-Milwaukee (May 2014).

TABLE I: Convergence of the total energies E (in *a.u.*) determined for the ground 1^1S -state of the Mu^- ion. The notation N is the total number of basis functions used.

N	E (series A)	E (series B)
400 ^(a)	-0.525054806501688	-0.525054806501688
3300	-0.525054806501730774024348	-0.52505480650173077402418
3500	-0.525054806501730774025187	-0.52505480650173077402500
3700	-0.525054806501730774025663	-0.52505480650173077402547
3840	-0.525054806501730774025914	-0.52505480650173077402573

^(a)The short-term cluster wave function with the carefully optimized non-linear parameters.

TABLE II: The expectation values of some propeties (in atomic units $a.u.$) for the Mu^- and ${}^\infty\text{H}^-$ ions.

ion	$\langle r_{eN}^{-2} \rangle$	$\langle r_{ee}^{-2} \rangle$	$\langle r_{eN}^{-1} \rangle$	$\langle r_{ee}^{-1} \rangle$
Mu^-	1.10551221416860383	0.15325557249058329	0.6796545010765918638	0.309199389149722180
${}^\infty\text{H}^-$	1.11666282452542572	0.15510415256242466	0.6832617676515272224	0.311021502214300052
	$\langle r_{eN} \rangle$	$\langle r_{ee} \rangle$	$\langle r_{eN}^2 \rangle$	$\langle r_{ee}^2 \rangle$
Mu^-	2.7271829808562793277	4.4392800894508588984	12.074193967756014098	25.51453633212122043
${}^\infty\text{H}^-$	2.7101782784444203653	4.4126944979917277211	11.913699678051262274	25.202025291240331897
	$\langle r_{eN}^3 \rangle$	$\langle r_{ee}^3 \rangle$	$\langle r_{eN}^4 \rangle$	$\langle r_{ee}^4 \rangle$
Mu^-	77.63368957070895012	184.07731257386858773	663.917821205619807	1632.23450379046995
${}^\infty\text{H}^-$	76.02309704902717911	180.60560023017477483	645.144542412219375	1590.09460393948530
	$\langle r_{eN}^6 \rangle$	$\langle r_{ee}^6 \rangle$	$\langle r_{eN}^8 \rangle$	$\langle r_{ee}^8 \rangle$
Mu^-	$9.1202102538542393 \cdot 10^4$	$2.2148515807730636 \cdot 10^5$	$2.33859300715081 \cdot 10^7$	$5.527178579001105 \cdot 10^7$
${}^\infty\text{H}^-$	$8.7266142406959270 \cdot 10^4$	$2.1253344237081047 \cdot 10^5$	$2.20357186954914 \cdot 10^7$	$5.221867644754489 \cdot 10^7$
	$\langle [r_{32}r_{31}]^{-1} \rangle$	$\langle [r_{eN}r_{ee}]^{-1} \rangle$	$\langle [r_{32}r_{31}r_{21}]^{-1} \rangle$	$\langle \delta(\mathbf{r}_{eeN}) \rangle$
Mu^-	0.37822708480340377	0.25017716867248448	0.19728255748264837	$4.877792625 \cdot 10^{-3}$
${}^\infty\text{H}^-$	0.38262789034020545	0.25307756706456687	0.20082343962918944	$5.129778775 \cdot 10^{-3}$
	$\langle \delta(\mathbf{r}_{eN}) \rangle$	$\nu_{eN}^{(a)}$	$\langle \delta(\mathbf{r}_{eN}) \rangle$	$\nu_{ee}^{(a)}$
Mu^-	0.1621506817228056	-0.995186945189023	$2.681680526352034 \cdot 10^{-3}$	0.499999989504
${}^\infty\text{H}^-$	0.1645528728473590	-1.00000000001778	$2.737992126104611 \cdot 10^{-3}$	0.500000002446
	τ_{eN}	τ_{ee}	$\langle f \rangle$	$\langle \mathbf{r}_{31} \cdot \mathbf{r}_{32} / r_{31}^3 \rangle$
Mu^-	0.6492013693363036596	-0.1038138782222772366	0.048647215112582520641	-0.4597494719447642817
${}^\infty\text{H}^-$	0.6498715811920881669	-0.1051476935659779011	0.048648867204549608205	-0.4642618530806317014
	$\langle -\frac{1}{2} \nabla_e^2 \rangle$	$\langle -\frac{1}{2} \nabla_N^2 \rangle$	$\langle \nabla_e \cdot \nabla_e \rangle$	$\langle \nabla_e \cdot \nabla_N \rangle$
Mu^-	0.2611868445353802015	0.554370044808132862	0.0319963557373724590	-0.554370044808132862
${}^\infty\text{H}^-$	0.2638755082721885983	0.560630798396681918	0.0328797818523047217	-0.560630798396681918

^(a)The expected cusp values (in $a.u.$) for the ${}^\infty\text{H}^-$ ion are $\nu_{eN} = -1.0$ and $\nu_{ee} = 0.5$. For the Mu^- ion these expected cusp values (in $a.u.$) are $\nu_{eN} = -0.9951869451890226269841682$ (with our muon mass) and $\nu_{ee} = 0.5$.

TABLE III: The expectation values of some bound state properties in atomic units of the MuPs system ($\mu^+e_2^-e^+$). The notation μ designates the positively charged muon, while the notations e^- and e^+ denote the electron and positron, respectively. N is total number of basis functions used.

N	E	$\langle r_{\mu-e^+}^{-2} \rangle$	$\langle r_{\mu-e^+}^{-1} \rangle$	$\langle r_{\mu-e^+} \rangle$	$\langle r_{\mu-e^+}^2 \rangle$	$\langle r_{\mu-e^+}^3 \rangle$	$\langle r_{\mu-e^+}^4 \rangle$
1000	-0.78631714258	0.1708472	0.3460478	3.678250	16.41087	86.4025	527.328
1200	-0.78631714533	0.1708472	0.3460478	3.678250	16.41087	86.4025	527.325
N	$\langle \frac{1}{2}p_{e^-}^2 \rangle$	$\langle r_{e^--e^+}^{-2} \rangle$	$\langle r_{e^--e^+}^{-1} \rangle$	$\langle r_{e^--e^+} \rangle$	$\langle r_{e^--e^+}^2 \rangle$	$\langle r_{e^--e^+}^3 \rangle$	$\langle r_{e^--e^+}^4 \rangle$
1000	0.323384673	0.3485151	0.4179003	3.488261	15.66631	85.1148	539.825
1200	0.323384675	0.3485152	0.4179000	3.488261	15.66630	85.1148	539.825
N	$\langle \frac{1}{2}p_{e^+}^2 \rangle$	$\langle r_{e^--\mu}^{-2} \rangle$	$\langle r_{e^--\mu}^{-1} \rangle$	$\langle r_{e^--\mu} \rangle$	$\langle r_{e^--\mu}^2 \rangle$	$\langle r_{e^--\mu}^3 \rangle$	$\langle r_{e^--\mu}^4 \rangle$
1000	0.136686711	1.19458809	0.7257295	2.326014	7.917324	35.9569	204.744
1200	0.136686710	1.19458811	0.7257295	2.326014	7.917324	35.9569	204.744
N	$\langle \frac{1}{2}p_{\mu}^2 \rangle$	$\langle r_{e^--e^-}^{-2} \rangle$	$\langle r_{e^--e^-}^{-1} \rangle$	$\langle r_{e^--e^-} \rangle$	$\langle r_{e^--e^-}^2 \rangle$	$\langle r_{e^--e^-}^3 \rangle$	$\langle r_{e^--e^-}^4 \rangle$
1000	0.591695472	0.2115948	0.36857646	3.594566	16.05697	86.0567	541.012
1200	0.591695470	0.2115948	0.36857646	3.594566	16.05697	86.0567	541.012
N	$\langle \delta(\mathbf{r}_{e^--e^+}) \rangle^{(a)}$	$\langle \delta(\mathbf{r}_{\mu-e^+}) \rangle^{(a)}$	$\langle \delta(\mathbf{r}_{\mu-e^-}) \rangle^{(a)}$	$\langle \delta(\mathbf{r}_{e^--e^-}) \rangle$	$\langle \delta(\mathbf{r}_{\mu e^-e^+}) \rangle$	$\langle \delta(\mathbf{r}_{e^--e^-e^+}) \rangle$	$\langle \delta(\mathbf{r}_{\mu e^+e^-e^-}) \rangle$
1000	0.02441854	0.00162037	0.17431747	0.00430157	$8.53207 \cdot 10^{-4}$	$3.66758 \cdot 10^{-4}$	$1.8176 \cdot 10^{-4}$
1200	0.02441854	0.00162037	0.17431728	0.00430159	$8.53185 \cdot 10^{-4}$	$3.67647 \cdot 10^{-4}$	$1.8176 \cdot 10^{-4}$

^(a)In the main text these expectation values are designated by the notations $\langle \delta(\mathbf{r}_{+-}) \rangle$, $\langle \delta(\mathbf{r}_{+\mu}) \rangle$ and $\langle \delta(\mathbf{r}_{-\mu}) \rangle$, respectively.

TABLE IV: The expectation values of a number of bound state properties (in atomic units) of the $p\mu e_2$, $d\mu e_2$ and $t\mu e_2$ four-body systems. The notation μ designates the positively charged muon, while the notation a stands for the protium, deuterium and tritium, respectively. The notation e^- denotes the electron.

system	E	$\langle r_{\mu-e}^{-2} \rangle$	$\langle r_{\mu-e}^{-1} \rangle$	$\langle r_{\mu-e} \rangle$	$\langle r_{\mu-e}^2 \rangle$	$\langle r_{\mu-e}^3 \rangle$	$\langle r_{\mu-e}^4 \rangle$
$p\mu e_2$	-1.150054535	1.531686	0.886755	1.609459	3.29137	8.1019	23.300
$d\mu e_2$	-1.150704304	1.532542	0.887313	1.607818	3.28330	8.0659	23.131
$t\mu e_2$	-1.150883319	1.532220	0.887597	1.607324	3.28200	8.0655	23.151
N	$\langle \frac{1}{2}p_{e^-}^2 \rangle$	$\langle r_{e^-a^+}^{-2} \rangle$	$\langle r_{e^-a^+}^{-1} \rangle$	$\langle r_{e^-a^+} \rangle$	$\langle r_{e^-a^+}^2 \rangle$	$\langle r_{e^-a^+}^3 \rangle$	$\langle r_{e^-a^+}^4 \rangle$
$p\mu e_2$	0.568945309	1.543128	0.889721	1.604998	3.27416	8.0409	23.071
$d\mu e_2$	0.569517734	1.544663	0.890487	1.603101	3.26538	8.0038	22.904
$t\mu e_2$	0.569758745	1.545693	0.890851	1.602424	3.26298	7.9972	22.892
N	$\langle \frac{1}{2}p_{\mu}^2 \rangle$	$\langle r_{a^+-\mu}^{-2} \rangle$	$\langle r_{a^+-\mu}^{-1} \rangle$	$\langle r_{a^+-\mu} \rangle$	$\langle r_{a^+-\mu}^2 \rangle$	$\langle r_{a^+-\mu}^3 \rangle$	$\langle r_{a^+-\mu}^4 \rangle$
$p\mu e_2$	2.408514781	0.480664	0.682547	1.509747	2.34618	3.7491	6.154
$d\mu e_2$	2.455562210	0.481420	0.683319	1.507081	2.33651	3.7228	6.090
$t\mu e_2$	2.468316240	0.482131	0.683880	1.505687	2.33209	3.7122	0.607
N	$\langle \frac{1}{2}p_{a^+}^2 \rangle$	$\langle r_{e^-e^-}^{-2} \rangle$	$\langle r_{e^-e^-}^{-1} \rangle$	$\langle r_{e^-e^-} \rangle$	$\langle r_{e^-e^-}^2 \rangle$	$\langle r_{e^-e^-}^3 \rangle$	$\langle r_{e^-e^-}^4 \rangle$
$p\mu e_2$	2.428621303	0.488805	0.569490	2.241373	6.01955	18.7208	65.961
$d\mu e_2$	2.477319166	0.489587	0.569990	2.239114	6.00633	18.6525	65.601
$t\mu e_2$	2.490270302	0.490284	0.570225	2.238367	6.00299	18.6410	65.570
N	$\langle \delta(\mathbf{r}_{e^--\mu}) \rangle$	$\langle \delta(\mathbf{r}_{\mu-a^+}) \rangle$	$\langle \delta(\mathbf{r}_{a^+-e^-}) \rangle$	$\langle \delta(\mathbf{r}_{e^-e^-}) \rangle$	$\langle \delta(\mathbf{r}_{a^+e^-e^-}) \rangle$	$\langle \delta(\mathbf{r}_{\mu e^-e^-}) \rangle$	$\langle \delta(\mathbf{r}_{\mu a^+e^-}) \rangle$
$p\mu e_2$	0.21307	$4.7078 \cdot 10^{-6}$	0.21489	0.01598	0.017084	0.017109	$2.393 \cdot 10^{-5}$
$d\mu e_2$	0.21139	$5.3055 \cdot 10^{-6}$	0.21318	0.01604	0.016533	0.016479	$2.233 \cdot 10^{-5}$
$t\mu e_2$	0.20881	$6.9770 \cdot 10^{-6}$	0.21358	0.01647	0.016308	0.016300	$2.157 \cdot 10^{-5}$